

University of Kentucky, Physics 306
Homework #1, Rev. A, due Wednesday, 2024-01-17

1. Perform the following exercises in **Mathematica**. The **Typewriter** font indicates command names.

- a) Plot `Exp[-1/x]` and its Derivatives—it transitions very smoothly to 0 for all $x < 0$.
- b) Find the `Series` of $(1+x)^a$ about $x=0$ to deduce the binomial expansion.
- c) Build a Plot of $x/(a^2 - x^2)$ piece by piece, starting from x^2 , then $a^2 - x^2$, etc.
- d) Plot the functions `Sin[x]`, `Sin[x-a]`, `Sin [x/b]`, `Sin[(x-a)/b]`, `Sin[x]^2`, `ArcSin[x]` using your own numbers for constants a and b .
- e) Show graphically that i) $\cos^2(x) = (1 + \cos(2x))/2$, ii) $\sin^2(x) = (1 - \cos(2x))/2$, iii) $2\cos(x)\sin(x) = \sin(2x)$, iv) $\cos^2(x) + \sin^2(x) = 1$, v) $(\cos(x) - \sin(x))(\cos(x) + \sin(x)) = \cos(2x)$.
- f) Manipulate a Plot of $f(t) = x_0 \text{Exp}[-\alpha t] \text{Sin}[\omega t]$ with sliders for the parameters x_0 , α , ω to investigate their effect on damped oscillatory motion, and also $f(x,t) = \text{Cos}[x - 2 \text{Pi } t]$ versus x with sliding parameter t to see the wave travel.
- g) Solve $x^2 + 1 = 0$ for x and substitute $x \rightarrow I$ into the LHS $x^2 + 1$ using the `/.` operator.
- h) Show Euler's theorem: `Exp[I x]==Cos[x] + I Sin[x]` by `Series` expansion of both sides. [*bonus*: Solve for $\cos(x)$ and $\sin(x)$ in terms of e^{ix} and e^{-ix} .]
- i) Plot the functions `Exp[x]/2`, `-Exp[x]/2`, `Exp[-x]/2`, `-Exp[-x]/2` along with the hyperbolic functions `Sinh[x]`, `Cosh[x]`, `-Sinh[x]`, `-Cosh[x]` and compare your results with h).
- j) Show that `Cosh[x]^2-Sinh[x]^2 = 1` and compare your results with e). [*bonus*: Parametrically plot $(x,y) = \{\text{Cos}[t], \text{Sin}[t]\}$ and compare with $\{\text{Cosh}[t], \text{Sinh}[t]\}$. What is the relation to the above identities? Thus they are called circular (elliptical) and hyperbolic functions.]
- k) Show graphically that `ArcSinh[y] = Log[y+Sqrt[y^2+1]]`. [*bonus*: What is `Tanh` in terms of `Exp`? Compare the *sigmoid* functions `Tanh[x]` and `2/Pi ArcTan[Pi/2 x]`.]
- l) [*bonus*: Explore plots of the matrix equation $\{x,y\} \cdot \{\{a,b\}, \{b,c\}\} \cdot \{x,y\} = 0$ for various values of (a,b,c) to discover conic sections. Calculate `Det[{a,b},{b,c}]` for each.]
- m) [*bonus*: Normalize the normal distribution $p(x) = \text{Exp}[-((x-\mu)/\text{sig})^2/2]$ so that it Integrates to 1 over the real line $-\infty < x < \infty$. Plot the normal distribution and the cumulative distribution $P(x) = \int_{-\infty}^x p(x)$ for $\mu = 2$ and $\sigma = 3$.]
- n) [*bonus*: Show that the three functions $f(x,y) = \text{Cos}[x] \{\text{Exp}[-y], \text{Sinh}[y], \text{Cosh}[y]\}$ are all solutions of $\nabla^2 f = 0$. `ContourPlot` the functions.]
- o) [*bonus*: `ContourPlot` the Re, Im parts of $f(z) = z^2 = (x + I y)^2$, `Cos[z]`, `Sin[z]`, `Exp[z]`.]

2. Taylor Series approximations allow focusing on the essential physics and analytically calculating problems with no closed form solution.

a) Calculate the Taylor series $f(x) = f(a) + f'(a)(x-a) + \frac{1}{2}f''(a)(x-a)^2 + \frac{1}{3!}f^{(3)}(a)(x-a)^3 + \dots$ of $f(x) = e^x$, $\cos(x)$, $\sin(x)$ to 5th order (x^5) about $a = 0$ and compare with Mathematica.


b) Perform long division of $\tan(x) = \sin(x)/\cos(x)$ from the series above to obtain the expansion for $\tan(x)$ to 5th order in x and compare with Mathematica.

c) Use the binomial series $(1+x)^\alpha = 1 + \frac{\alpha}{1}x + \frac{(\alpha)(\alpha-1)}{(1)(2)}x^2 + \dots$ to obtain the **multipole expansion** of the electrostatic potential $\vec{V}(\vec{r}) = q/4\pi\epsilon z$, where $z = ((\vec{r} - \vec{r}') \cdot (\vec{r} - \vec{r}') = r^2 + 2rr' \cos \gamma + r'^2)^{1/2}$ to 2nd order in r' . [*bonus*: Identify the monopole, dipole, and quadrupole moments and potentials.]

d) In the small-area approximation, the **solid angle** subtended by a detector of area $|\vec{A}|$ with unit normal \hat{A} located at spherical coordinates (R, Θ, Φ) , tilted down by the angle Γ (from facing the origin) is

$$\Omega(x, y) = \frac{\vec{A} \cdot \hat{D}}{D^2} = \frac{A}{R^2} \frac{\cos \Gamma - \sin(\Theta + \Gamma)(x \cos \Phi + y \sin \Phi)}{(1 - 2 \sin \Theta (x \cos \Phi + y \sin \Phi) + x^2 + y^2)^{3/2}}.$$

Expand $\Omega(x, y)$ to second order in x and y about the origin. [*bonus*: Show that for the total solid angle of 4 symmetric detectors at azimuthal angles $\Phi = 0, \pi/2, \pi, 3\pi/2$, all terms vanish except $\Omega(x, y) = \Omega_0 + \Omega_2(x^2 + y^2)$. Calculate the relation between Θ and Γ that forces $\Omega_2 = 0$]



A view of two detectors centered at $(R, \Theta, 0)$ and $(R, \Theta, \pi/2)$ in spherical coordinates.