

L24-Postulates of QM

Wednesday, November 18, 2015 10:24

* what do you do in:

Classical mechanics? predict future measurements:

calculate (evolve) trajectory, (use conservation principles)

Quantum mechanics? predict what happens to a state

a) how it evolves in time undisturbed

b) what happens when you make a measurement

- note: a QM state is DEFINED by what you can measure!

* Practically, what do we do to solve problems in QM?

a) calculate eigenfunctions of operators (Hermitian)

b) rotate vectors or change basis (Unitary)

- solve $\hat{H}|\psi_n\rangle = E_n|\psi_n\rangle$ to get stationary states.

- change basis of initial state to $|\Phi_0\rangle = \sum c_n |\psi_n\rangle$ energy basis

- rotate stationary states in time: $|\Psi(t)\rangle = \sum c_n |\psi_n\rangle e^{iE_n t/\hbar}$

- solve eigenfunctions of an observable $\hat{Q}|\phi_n\rangle = q_n|\phi_n\rangle$

- rotate basis to observable states: $|\Psi(t)\rangle = \sum a_n(t) |\phi_n\rangle$

to calculate probability $|a_n|^2$ of measuring q_n

- project state (collapse wavefn) $|\Psi\rangle \rightarrow |\phi_n\rangle$ after measuring q_n

What are the tools? inner product $\langle\psi|\phi\rangle$

- orthonormality: $\langle\phi_m|\phi_n\rangle = \delta_{nm}$ closure: $\sum_n |\phi_n\rangle\langle\phi_n| = I$

* Postulates of QM:

1) Hilbert space of states: superposition postulate

state $|\psi\rangle$ = collection of complex probability amplitudes $\psi(x)$ or c_n

(vector components) which linearly combine to form new states
 It has an inner product $\langle \Psi | \Psi \rangle = \int dx |\Psi(x)|^2 = \sum_n |c_n|^2$
 and is normalizable $\langle \Psi | \Psi \rangle = 1$ so probabilities add to 100%

- 2) Hermitian observables: expansion/projection postulate
 observable = collection of determinate states & measurements
 (operator $\hat{Q}$) with real eigenvalues q_n & orthogonal eigenvectors $|\phi_n\rangle$
 which form a complete set of basis vectors: $|\Psi\rangle = \sum_n a_n |\phi_n\rangle$
 - $|a_n|^2$ = probability of measuring q_n
 - $|\Psi\rangle \rightarrow |\phi_n\rangle$ after measuring q_n

- 3) Hamiltonian: evolution postulate
 states evolve in time according to Schrödinger Eq: $\hat{H}|\Psi\rangle = \hat{E}|\Psi\rangle$
 - eigenstates of $\hat{H}$ are stationary states: $\hat{H}|\psi_n\rangle = E_n |\psi_n\rangle$
 - evolution of mixed states: $|\Psi(x,t)\rangle = \sum_n c_n |\psi_n\rangle e^{-iE_n t/\hbar}$

- 4) Heisenberg: uncertainty postulate
 - canonical commutation relation $[\hat{x}, \hat{p}] = i\hbar$ for conjugate observables
 - position and momentum are complementary: $\Delta x \Delta p \geq \hbar/2$
 - momentum operator $\hat{p} = -i\hbar \frac{\partial}{\partial x}$ and eigenstates $\psi_p(x) = e^{ipx/\hbar}$

- 5) Pauli: exclusion postulate
 - identical particle exchange symmetry $\Psi(x_1, x_2) = \pm \Psi(x_2, x_1)$
 - only one fermion can occupy each state
 (will discuss next semester)