

**University of Kentucky, Physics 520**  
**Homework #9, Rev. A, due Wednesday, 2016-11-08**

**0.** Griffiths [2ed] App. A #4, #9, #10.

**1.** The **complex plane**  $\{\mathbf{w} = (x, y)\}$  is the vector space of *real* and *imaginary* components of  $w = x + iy \in \mathbb{C}$ , with the additional operation of multiplication, similar to the *general linear group*  $GL(n)$  of  $n \times n$  matrices. We explore this analogy by generalizing the exponential map  $e^{i\phi}$ .

**a)** Identify the two basis elements of the complex plane in the above representation of  $w$ .

**b)** Show that the dot and cross product of two points  $\mathbf{w}_1 = (x_1, y_1)$  and  $\mathbf{w}_2 = (x_2, y_2)$  are given by the real and imaginary parts of the complex product  $w_1^* w_2 = \mathbf{w}_1 \cdot \mathbf{w}_2 + i(\mathbf{w}_1 \times \mathbf{w}_2)_z$ , where  $w^* = x - iy$  is the *complex conjugate* of  $w$ . Identify the symmetric and antisymmetric terms of this product. Thus the complex product  $|w|^2 = w^* w$  equals the vector product  $\mathbf{w} \cdot \mathbf{w}$ .

**c)** Show graphically that the operator  $w \rightarrow iw$  rotates the point  $w$   $90^\circ$  CCW about the origin.

**d)** Show graphically that the operator  $1 + i d\phi : w \mapsto w + iw d\phi$  preserves the magnitude of  $w$  (assuming  $d\phi^2 = 0$ ), but rotates it CCW by the infinitesimal angle  $d\phi$ .

**e)** Obtain a finite rotation from an infinite number of  $d\phi$  rotations as follows: formally integrate the equation  $dw = iw d\phi$  with the initial condition  $w|_{\phi=0} = w_0$  to obtain the rotation formula  $w(\phi) = R_\phi w_0$ , where  $R_\phi = e^{i\phi}$ . Use this result to justify the identity  $e^x = \lim_{n \rightarrow \infty} (1 + \frac{x}{n})^n$ .

**f)** Separate the Taylor expansion of  $e^{i\phi}$  into  $x+iy$  to prove Euler's formula,  $e^{i\phi} = \cos \phi + i \sin \phi$ .

**g)** Show that complex multiplication by  $i$  is equivalent to the vector operator  $\hat{\mathbf{z}} \times$ .

**h)** Determine the matrix representation  $M_z$  of the operator  $\hat{\mathbf{z}} \times$ , where  $M_z \mathbf{r} = \hat{\mathbf{z}} \times \mathbf{r}$ . Do the same for  $M_x$  and  $M_y$  to show that  $\mathbf{v} \times = \mathbf{v} \cdot \mathbf{M} = v_x M_x + v_y M_y + v_z M_z$  is the matrix representation of  $\mathbf{v} \times$  for any vector  $\mathbf{v}$ . *Hint:* You should find that the vector of matrices  $\mathbf{M} \sim (M_i)_{jk} = \varepsilon_{ijk}$  (cross product tensor) is completely antisymmetric in indices  $i, j, k$ .

**i)** Restricting to the  $xy$ -plane, show that the  $2 \times 2$  matrix  $M_z^2 = -I$  analogous to  $i^2 = -1$ , and the matrix for a CCW rotation  $\phi$  is  $R_\phi = e^{M_z \phi} = I \cos \phi + M_z \sin \phi = \begin{pmatrix} \cos \phi & -\sin \phi \\ \sin \phi & \cos \phi \end{pmatrix}$ . *Hint:* the *exponential* of a matrix  $M_z$  is defined by its Taylor expansion as in part f). Note that in general, the matrix  $R_{\mathbf{v}}$  for a CCW rotation by angle  $v = |\mathbf{v}|$  about the  $\hat{\mathbf{v}}$ -axis can be written as  $R_{\mathbf{v}} = I \cos v + \mathbf{M} \cdot \hat{\mathbf{v}} \sin v + \hat{\mathbf{v}} \hat{\mathbf{v}}^T (1 - \cos v)$ , where the third term corrects for the non-rotating projection along  $\mathbf{v}$ .

**j)** Calculate the eigenvalues and eigenvectors of  $M_z = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}$  to show that  $M_z = VWV^\dagger = \frac{1}{2} \begin{pmatrix} 1 & 1 \\ i & -i \end{pmatrix} \begin{pmatrix} -i & 0 \\ 0 & i \end{pmatrix} \begin{pmatrix} 1 & -i \\ 1 & i \end{pmatrix}$  and  $e^{M_z \phi} = V e^{W \phi} V^\dagger = \frac{1}{2} \begin{pmatrix} 1 & 1 \\ i & -i \end{pmatrix} \begin{pmatrix} e^{-i\phi} & 0 \\ 0 & e^{i\phi} \end{pmatrix} \begin{pmatrix} 1 & -i \\ 1 & i \end{pmatrix}$ . Multiply this out to verify part i). Thus real Hermitian matrices have real eigenvalues while antiHermitian matrices have  $\text{Tr}=0$  and imaginary eigenvalues. The exponential of a Hermitian matrix is positive definite with real positive eigenvalues, while the exponential of an antiHermitian matrix is unitary with  $\text{Det}=1$  and unit modulus eigenvalues. This is the normal matrix analogy: Hermitian/antiHermitian matrices are analogous to the real/imaginary numbers.