University of Kentucky, Physics 521 Homework #15, Rev. B, due Monday, 2016-04-25

0. Griffiths [2ed] Ch. 9 #9.1, #9.2, #9.6, #9.7, #9.8, #9.9.

1. The Lyman-alpha (L_{α}) transition $(n = 2 \rightarrow 1)$ is the highest energy, shortest wavelength line of the hydrogen spectrum.

- a) Calculate the energy and wavelength of this transition.
- b) Which $n = 2 \rightarrow 1$ substates satisfy the electric dipole selection rules?
- c) Calculate the lifetime and line width of this transition.

d) Calculate the effects of collisional broadening [use $\sigma_{col} = \pi(\langle r^2 \rangle_{100} + \langle r^2 \rangle_{211})$, see Griffiths problem #4.13], Doppler broadening, and recoil line shift at STP (20°C, 1 atm), as fractions of the natural wavelength and line width. [Note, however, that UV light attenuates rapidy in air!]

2. Lasers require a population inversion of occupation numbers $N_b > N_a$ to achieve a higher rate of stimulated emission than of absorption of photons in the transition $E_b \rightarrow E_a$. An optical cavity with mirrors at each end reflects the laser beam back and forth to create a standing wave, enhancing the photon density and narrowing the line width. A partial silvered mirror on one end allows a fraction of the photons to exit as the laser beam, contributing to attenuation of photons in the cavity. Lasing occurs if there is a net gain of photons in the laser due to stimulated emission.

a) Integrate the spectral density of the Breit-Wigner resonance

$$\rho(\omega) = \rho(\omega_0) \frac{(\Gamma/2)^2}{(\omega - \omega_0)^2 + (\Gamma/2)^2}$$

to obtain the energy density $u = \int_0^\infty \rho(\omega) d\omega \approx \int_{-\infty}^\infty \rho(\omega) d\omega$ in the cavity. Convert this to N_γ , the number of photons in the beam, which occupies a volume V in the cavity.

b) Show that N_{γ} , the rate of change of the number of photons in the cavity [due to stimulated emission, absorption, and attenuation] is positive if the population inversion is greater than the *critical density*

$$\Delta n_c \equiv \frac{N_b - N_a}{V} = \frac{\omega^2 \ \Gamma \tau_s}{2\pi c^3 \tau_\gamma} = \frac{2\pi \Gamma \tau_s}{\lambda^2 c \tau_\gamma},$$

neglecting the degeneracy of states, where τ_s is the lifetime of spontaneous emission, and τ_{γ} is the characteristic lifetime of photons in the laser cavity.

c) Calculate the population inversion density needed for a Ruby laser [the first pulsed laser] and a He-Ne laser [the first CW laser], using the following values [Tipler & Llewellyn, 5ed]:

Ruby laser He-Ne laser

λ	$694.3 \mathrm{~nm}$	632.8 nm
n (refractive index)	1.76	1.00
$ au_s$	$3 \mathrm{\ ms}$	100 ns
$ au_\gamma$	29 ns	330 ns
$\delta u = \Gamma/2\pi$	$330 \mathrm{GHz}$	0.9 GHz.

3. [bonus] Design an efficient, economic Lyman-alpha laser.