

- * Final Exam - cumulative from whole semester
 - must solve 4/5 problems (see midterm)
 - each with a calculation & conceptual questions

4) Spin & Angular momentum

Quantum numbers & ladder operators.

Pauli Matrices, diagonalization,
combination of 2 spin- $\frac{1}{2}$ systems;
evolution of spin system in B-field.

5) Identical particles

Symmetrization of multi particle wave fn: $\psi_{\text{even}}, \psi_{\text{odd}}$.

Atomic configuration

Statistical counting of microstates from single-particle states.

Description of M-B, F-D, BE distributions. , Bose condensation.

Blackbody spectrum

6) Time-independent perturbation theory

solve nondeg. & degenerate first-order energy & eigenstates
using either operator wave functions or perturbation matrix

[hyper]fine structure: show magnetic interaction produces $\mathbf{L} \cdot \mathbf{S}$ coupling
identify states with good $\mathbf{L} \cdot \mathbf{S}$, $\mathbf{J} \cdot \mathbf{S}$

Zeeman effect: states & energies in low/strong field limit.

7) Variational Principle

problem to minimize energy of parameterized state.

9) Time dependent perturbation theory:

calculate evolution $[c_a(t), c_b(t)]$ to first order

calculate a transition probability:

derive relation between A, B coefficients.

Explain Breit-Wigner line shape, selection rules.

11) Scattering

explain the relation between various waves, amplitudes, and cross sections

solve for partial wave scattering amplitude.

* Fermi's golden rule: connection between Born Approximation & time dependent perturbation theory

$$R_{i \rightarrow f} = \frac{2\pi}{\hbar} |\langle f | H' | i \rangle|^2 \rho(E') = \frac{\sigma \cdot V}{V} \quad \rho(E') = \frac{dn}{dE'} = \frac{V 4\pi p^2 dp d\Omega}{(2\pi\hbar)^3 dE'}$$

$$\sigma = \frac{2\pi}{\hbar V} |\langle f | H' | i \rangle|^2 \rho(E') V = \frac{V^2 E'^2}{(2\pi)^2 (\hbar c)^4} |\langle f | H' | i \rangle|^2 \Delta\Omega$$

- compare to rate of stimulated emission

$$R_{i \rightarrow f} = \frac{\pi}{3\epsilon_0 \hbar^2} \langle f | e \vec{r} | i \rangle^2 \rho(\omega_0) = \frac{\pi \hbar \omega^2 \rho(\omega_0)}{3\epsilon_0 \hbar^2}$$

- scattering is a transition from one state to another



$$f(\theta, \phi) = \frac{-m}{2\pi\hbar^2} \langle f | V | i \rangle = -\frac{m}{2\pi\hbar^2} \int d^3\vec{r} e^{i\vec{q} \cdot \vec{r}} V(\vec{r})$$

* Elastic scattering & form factors:


$$\begin{aligned} \langle f | H' | i \rangle &= \frac{1}{V} \langle e^{i\vec{k}' \cdot \vec{r}} | e \phi(\vec{r}) | e^{-i\vec{k} \cdot \vec{r}} \rangle = \int e \phi(\vec{r}) e^{i\vec{q} \cdot \vec{r}} d^3\vec{r} \\ &= \int e [\nabla^2 \rho(\vec{r}) / \epsilon] e^{i\vec{q} \cdot \vec{r}} d^3\vec{r} = \frac{e}{q^2 \epsilon} F(\vec{q}) \end{aligned}$$

$$q^2 = (\vec{p}' - \vec{p})^2 = p'^2 + p^2 - 2pp' \cos \theta = 4p^2 \sin^2 \theta/2 \quad [p=p']$$

$$\frac{d\sigma}{d\Omega} = \underbrace{\frac{Z^2 \alpha^2 (\hbar c)^2}{4E^2 \sin^4 \theta/2}}_{\text{Rutherford cross. sec.}} \cdot \underbrace{|F(q)|^2}_{\text{form factor}}$$



QED: $\langle f | H' | i \rangle \Rightarrow$ invariant amplitude $M = J_\mu \frac{g_{\mu\nu}}{q^2} J_\nu$
 massless photon γ has propagator $\frac{g_{\mu\nu}}{q^2}$
 connecting 2 scattering currents:
 $J^\mu = \langle f | e \gamma^\mu | i \rangle \quad \gamma^\mu \sim \text{spin.}$



Impulse expansion: $G(q) = \frac{1}{q^2 + k^2} \sim \frac{e^{ikr}}{r}$ propagator for T.S.E.

Massive particles: $G(q) = \frac{1}{q^2 - m^2} \sim \frac{e^{-mr}}{r}$ Yukawa potential