

**University of Kentucky, Physics 521**  
**Homework #18, Rev. A, due Monday, 2017-04-17**

**0.** Griffiths [2ed] Ch. 9 #1, #6, #7, #8, #9.

**1.** The **Lyman-alpha** ( $L_\alpha$ ) line ( $n = 2 \rightarrow 1$ ) is the lowest energy transition to the atomic hydrogen ground state. It is an important wavelength for spectroscopy and laser-cooling antihydrogen.

a) Calculate the energy and wavelength of this transition.

b) Which  $n = 2 \rightarrow 1$  substates satisfy the electric dipole selection rules?

c) Calculate the lifetime and line width of this transition.

d) Calculate the effects of collisional broadening [use  $\sigma_{col} = \pi(\langle r^2 \rangle_{100} + \langle r^2 \rangle_{211})$ , see Griffiths problem #4.13], Doppler broadening, and recoil line shift at STP (20°C, 1 atm), as fractions of the natural wavelength and line width. [Note, however, that UV light attenuates rapidly in air!]

**2. Lasers** require a *population inversion* of occupation numbers  $N_b > N_a$  to achieve a higher rate of *stimulated emission* than of absorption of photons in the transition  $E_b \rightarrow E_a$ . An *optical cavity* with mirrors at each end reflects the laser beam back and forth to create a standing wave, enhancing the photon density and narrowing the line width. A partial silvered mirror on one end allows a fraction of the photons to exit as the laser beam, contributing to attenuation of photons in the cavity. Lasing occurs if there is a net gain of photons in the laser due to stimulated emission.

a) Integrate the spectral density of the Breit-Wigner resonance

$$\rho(\omega) = \frac{\rho(\omega_0)(\Gamma/2)^2}{(\omega - \omega_0)^2 + (\Gamma/2)^2}$$

to obtain the energy density  $u = \int_0^\infty \rho(\omega) d\omega \approx \int_{-\infty}^\infty \rho(\omega) d\omega$  in the cavity. Convert  $u$  to  $N_\gamma$ , the number of photons in the beam, which occupies a volume  $V$  in the cavity.

b) Show that  $\dot{N}_\gamma$ , the rate of change of the number of photons in the cavity [due to stimulated emission, absorption, and attenuation] is positive if the population inversion is greater than the *critical density*

$$\Delta n_c \equiv \frac{N_b - N_a}{V} = \frac{\omega^2 \Gamma \tau_s}{2\pi c^3 \tau_\gamma} = \frac{2\pi \Gamma \tau_s}{\lambda^2 c \tau_\gamma},$$

neglecting the degeneracy of states, where  $\tau_s$  is the lifetime of spontaneous emission, and  $\tau_\gamma$  is the characteristic lifetime of photons in the laser cavity.

c) Calculate the population inversion density needed for a Ruby laser [the first pulsed laser] and a He-Ne laser [the first CW laser], using the following values [Tipler & Llewellyn, 5ed]:

	Ruby laser	He-Ne laser
$\lambda$	694.3 nm	632.8 nm
$n$ (refractive index)	1.76	1.00
$\tau_s$	3 ms	100 ns
$\tau_\gamma$	29 ns	330 ns
$\delta\nu = \Gamma/2\pi$	330 GHz	0.9 GHz.

**3. [bonus]** Design an efficient, economic Lyman-alpha laser.